Children and young people with eating disorders

Guidelines for primary care professionals

These guidelines share advice about spotting the signs of eating disorders and when to refer children and young people to the community eating disorders service (CEDS) for specialist outpatient support.

Remember to refer ALL suspected children and young {<u>`</u>} people with eating disorders to your local CEDS

- History of weight loss (> 1kg a week for 2 consecutive weeks) or failure to gain weight
- If no weight available has clothes size changed?
- Have parents noticed weight loss?
- How long has the young person been trying to lose weight & what have they tried?

52

- If they are binging, vomiting, using laxatives, how often in a week?
- How much are they currently eating & drinking?

Children and young people DO NOT need to be exceptionally underweight for a referral

• 'Low weight' is now defined as a loss of 10% of a young person's expected body weight

When to refer

Refer ALL Children and Young People with a suspected eating disorder to your local CEDS

- Anxiety about gaining weight behaviours to lose weight that are more than a 'teenage diet' such as extreme restricting or frequent binging
- Low, normal or overweight with the above behaviours
- Fainting, dehydration and confusion

- imescales
- Urgent assessment will be within 5 days
- 15 days if routine NICE concordant treatment within 28 days (note this is primarily family based treatment rather than individual)

• Are they over exercising?

- If appropriate, when did they last have a menstrual cycle?
- Any risks?

20



Include the physical exam

CONSENT:

aware of this referral?

• Are both the young person & parent(s)

• Who has given consent to this referral?

- Weight and height
- BP and pulse (sitting and standing)
- Temperature
- Send for baseline blood tests at the same time as referral: FBC, U+E's, LFT, Glucose, ESR, TFT, bone profile, Calcium, Magnesium
- DO NOT DELAY REFERRAL WAITING FOR RESULTS!

RED FLAGS

Phone your local CYP CEDS immediately

- not eating and fluid refusal for more than 24 hours
- cardiovascular compromise and fainting
- BMI <13: under 0.2nd centile
- BP < 0.4th centile
- Pulse <40 bpm sitting and standing/postural drop
- Temperature <35°C
- Weight loss > 1kg a week for 2 consecutive weeks

Frimlev Clinical Commissioning Group



Berkshire West Clinical Commissioning Group

Eating disorders. Know the first signs?







Are they obsessive about food?

Flips
Is their behaviour changing?

Hips Do they have distorted beliefs about their body size?



Kips

Are they often tired

or struggling to

concentrate?

Nips Do they disappear to the toilet after meals? Skips Have they started exercising excessively?

Online support

BERKSHIRE EATING DISORDER SERVICES -

BERKSHIRE HEALTHCARE FOUNDATION TRUST

For an urgent referral telephone our HealthHub

Disorders" module of PPEP Care training at this link

line (9am-5pm), Mondays to Fridays

Visit beateatingdisorders.org.uk for information about eating disorders, and to access online support groups and one-to-one chat.

Use helpfinder.beateatingdisorders.org.uk to find services in your area.

If you suspect a young person within your care has an eating disorder,

you can make a referral yourself. Once the referral is received, it will

be passed to the CYPF Berkshire Eating Disorders service for triage.

If you're unsure whether to make a referral please call our duty

You can sign up for the "Supporting young people with Eating

@BeatED@BeatEDSupportf beat.eating.disorders

Beat is the UK's eating disorder charity. We are a champion, guide and friend for anyone affected by an eating disorder.



Help for young people

Youthline: 0808 801 0711

Email: fyp@ beateatingdisorders.org.uk

For more referral information: helpfinder.beateatingdisorders.org.uk/refer Berkshire version adapted from the original work of Healthy London Partnership and BEAT with thanks.